

## Age and social-related risk factors for age-related macular degeneration among patients attending eye care at Jinja Regional Referral Hospital. A cross-sectional study.

Joshua Tumwebaze\*, Taddeo Mulumba  
Ophthalmic Clinical Officers' Training School, Jinja

Page | 1

### Abstract

#### Background

Age-related macular degeneration causes irreversible blindness. There are two types of Age-related Macular Degeneration, that is to say, Wet macular degeneration and Dry macular degeneration. The study aims to assess aim of the study is to assess the Age and social-related Risk factors for age-related macular Degeneration.

#### Methodology

A Cross-sectional descriptive study design involving quantitative methods of data collection.

The study population comprised all patients receiving eye care services for Age-related Macular Degeneration conditions at the eye unit of Jinja Regional Referral Hospital for a period of 6 months from October 2022 to March 2023. Non-probability sampling was used to select 50 participants.

#### Results

The majority, 29/50 (58%) respondents were females, and 21/50 (42%) were males. 39/50 (78%) of the respondents lived in Rural areas, while a lower number, 11/50 (22%), lived in Urban areas. The majority, 25/50 (50%), were between 61-70 years, followed by respondents 14/50 (28%) who were between 51-60 years, followed by respondents 2/50 (4%) who were between 41-50 years, while the least were respondents 2/50 (4%). Most 24/50(48%) strongly agreed that smoking is a risk factor for age-related Macular Degeneration. 32/50 (64%) took alcohol, while 18/50 (36%) did not take alcohol.

#### Conclusion

The significant risk factors for Age-related Macular Degeneration were increasing age groups of 61-70 years and 51-60 years, smoking, and a low level of education.

#### Recommendations

The Government should sensitize people on the risk factors for age-related Macular degeneration. At the same time, the Government should set up more eye clinics to increase accessibility to eye care services.

**Keywords:** Macular Degeneration, Smoking, Irreversible blindness, Jinja Regional Referral Hospital.

**Submitted:** June 24, 2024 **Accepted:** November 01, 2025 **Published:** April 04, 2026

**Corresponding Author:** Joshua Tumwebaze

**Email:** [tumwebazejoshua77@gmail.com](mailto:tumwebazejoshua77@gmail.com)

Ophthalmic Clinical Officers' Training School

#### Background

Age-related macular degeneration causes irreversible blindness among people aged 50 years and above. There are two types of Age-related Macular Degeneration, that is to say, Wet macular degeneration and Dry macular degeneration. Age-related Macular Degeneration (AMD) is a progressive disease that preferentially affects the macular region of the retina (Pennington and DeAngelis, 2016). About 170 million individuals have been affected by AMD globally, thus being the third leading cause of vision loss in the world (Wong et al., 2014). In the United States, the prevalence of AMD reaches 11 million, similar to that of invasive cancers combined, leading to an annual \$4.6 billion direct healthcare cost (Xiayan et al., 2020). 200 million people worldwide are estimated to have AMD, and by 2040,

this number is projected to rise to 300 million. By 2050, 5.4 million Americans are anticipated to be affected by rising patterns similar to this one (Deng et al., 2022).

Dry AMD causes blurred or reduced central vision due to thinning of the macula. It can affect one or both eyes. At any stage of dry AMD can turn into wet AMD. In the wet AMD, or exudative form of Age-related Macular Degeneration (AMD or ARMD), pathologic choroidal neovascular membranes (CNVM) develop under the retina. The CNVM can leak fluid and blood and, if left untreated, ultimately cause a centrally blinding disciform scar. (F Ryan Prall, Andrew A Dahl, 2023). Patients with age-related Macular Degeneration may experience visual distortions like straight lines or surfaces appearing to be bent or wavy. Low light may cause visual challenges, and you may not be

able to recognize faces as easily as you have in the past. (Bamini Gopinath, 2016). You might even experience a visual hallucination called Charles Bonnet syndrome. However, other signs may include Colors may look dull, or it may be difficult to tell the difference between subtle shades of colors or textures. What at first glance appears to be clumsiness may actually be related to your eyesight. Blind spots in your central vision might make it difficult to read, sew, and do close work. The study aims to assess aim of the study is to assess the Age and social-related risk factors for age-related macular Degeneration.

## **METHODOLOGY**

### **Study design.**

A Cross-sectional descriptive study design involving quantitative methods of data collection.

### **Study Area**

The study area was the eye clinic of Jinja Regional Referral Hospital. The hospital is located in the South Eastern region of the country, in the central division of Jinja city, near the source of the Nile, which is located 80km east of Kampala, the capital city of Uganda. The hospital was founded in 1962 and has a bed capacity of 600. Jinja hospital is a regional referral hospital serving several clients/ patients across the Busoga region and other neighboring districts of Buikwe, Kayunga, and others. The clients are referred from other hospitals and health centers IV, while others are self-referred. Among the services provided are medical, surgical, orthopedic, eye care, private, gynecology, pediatrics, dental, ENT, Lab, X-Ray/Scan, Immunization, HIV testing and counseling, Reproductive health services, and Safe Male Circumcision, among others.

The hospital has 15 wards, which include surgical female/male, medical female/male, TB, Eye, Urology, Grade A, Annex, TB, Psychiatric, and Children's wards, Intensive Care Unit, postnatal, Gynecological, and Maternity wards.

### **Study Population**

The study population comprised all patients receiving eye care services for Age-related Macular Degeneration conditions at the eye unit of Jinja Regional Referral Hospital.

### **Time scope**

The study was carried out for a period of 6 months, that is October 2022 to March 2023

### **Sample size determination.**

The sample size was calculated using the method below, according to Mbuto Samuel (2004)

Sample size =  $\frac{DxH}{T}$  Where:

Where:

D -number of days that were available for data collection

H -number of hours taken per day

T -total time that was spent on each event.

D =30days

H =5hours

T =3minutes

N =50participants

### **Sampling technique**

The study was a non-probability sampling process, specifically a purposive sampling technique, where only patients with age-related Macular Degeneration participated in the study.

### **Sampling procedure**

Patients registered at the reception, their age, and sex were documented regardless of presenting ocular complaints. Whenever any ophthalmic clinical officer or ophthalmologist identified a case of Age-related Macular Degeneration during the course of history taking, examination, or treatment of a patient, he/she directed the patient to the principal investigator or any of the research assistants for inclusion into the study.

### **Data collection method**

The principal investigator used interviews as the method of data collection using an interview guide with well-structured questions.

All clinicians at the JRRH eye clinic were sensitized and requested to identify the participants with Age-related Macular Degeneration of eye conditions.

### **Data collection tool**

A questionnaire filled in by the researcher or a research assistant was used to collect data from the respondents.

### **Data Collection procedure**

The principal investigator started by creating rapport and explained the objectives of the study to the respondents and reassured them about confidentiality. The questions were read and interpreted for the respondents to understand. Responses were given in the local language and written in English by the principal investigator and his assistants for those who did not understand English. Data was collected for a period of 30 days, as at least 2 respondents, each taking 20-30 minutes, were interviewed daily until the required sample size was obtained. At the end, the respondents were thanked for their cooperation.

### **Pre-testing:**

The study tool was pre-tested on a few patients attending the eye clinic, and necessary adjustments are to be made to ensure validity and reliability. Patients who were included

in the study were asked to consent. Those in exclusion were below the age range and those who did not consent to participate in the study.

### Data Analysis and Presentation

The data collected was manually analyzed through tallying and, use of scientific calculators. The results were presented in the form of tables, Figures, and statements. A computer was used to type the report. Microsoft Excel was used to draw charts and graphs for a clear presentation of the findings.

### Ethical Consideration

On approval of the research proposal by the school, a letter of introduction was provided by the Principal of Ophthalmic Clinical Officers Training School, and then it was delivered to the hospital director, JRRH, to grant permission to undertake the study in the hospital. The researcher introduced himself to the in-charges of the eye clinic. He then introduced himself and the objectives to the respondents. The principle of confidentiality and anonymity was observed throughout the study.

### Results: Social Demographic

**Table 1 Shows Social Demographic Characteristic ( n=50 Respondents)**

Category	Frequency(n=50)	Percentage (%)
<b>Age</b>		
35-40	02	04
41-50	09	18
51-60	14	28
61-70	25	50
<b>Sex</b>		
Female	29	58
Male	21	42
<b>Tribe</b>		
Musoga	27	54
Muganda	13	26
Munyankole	08	16
Others	02	04
<b>Religion</b>		
Anglican	19	38
Muslim	09	18
Catholic	22	44
<b>Level of Education</b>		
Primary	32	64
Secondary	16	32
Tertiary	02	04
<b>Occupation</b>		
Business personnel	10	20
House wife	19	38
Peasant	21	42
<b>Home Area</b>		
Rural	39	78
Urban	11	22

Table 1, on age, 25/50(50%), which is half of the total number of respondents, were between 61-70 years, and the minority 02/50 (04 %) were 31-40 years.

On sex, more respondents, 29/50 (58%) were females than 21/50 (42%) males.

On tribe, the majority 27/50 (54%) of the respondents were Basoga, while the minority 02/50 (04%) were among others.

On religion, most of the respondents, 22/50 (44%), were Catholics, and a minority, 09/50 (18%), were Muslims. The Level of Education established that the majority, 32/50 (64%) of the respondents were primary dropouts, while the least 02/50 (04%) had attained a tertiary school level of education.

On occupation, most of the respondents, 21/50 (42%), were peasants, while 10/50 (20%) were Business personnel. Concerning the home area, a higher number of 39/50 (78%) respondents lived in Rural areas, while a lower number, 11/50 (22%), lived in Urban areas.

### Age-related Risk factors for age-related macular Degeneration

**Figure 1 Age-related risk factors for age-related Macular degeneration**




Figure 1, the majority of the respondents, 25/50 (50%), were between 61-70 years, followed by respondents 14/50 (28%), who were between 51-60 years, followed by respondents 9/50 (18%) who were between 41-50 years, while the least were respondents 2/50 (4%).

### Social-related risk factors for age-related Macular Degeneration

**Table 2 Shows Smoking as a risk factor of age-related Macular Degeneration.**

SMOKING	NUMBER OF RESPONDENTS	PERCENTAGE (%)
Agree	20	40
Strongly Agree	24	48
Disagree	06	12

Table 2, The highest number of respondents 24/50(48%), strongly agreed that smoking is a risk factor for age-related Macular Degeneration, followed by 20/50(40%) respondents who agreed, while the least number of respondents, 06/50 (12%), Disagreed.

**Figure 2 A Pie Chart Showing Alcohol as a socially related risk factor for Age-Related Macular Degeneration.**




Figure 2, A larger number of respondents, 32/50 (64%), said YES, indicating that they took alcohol, than 18/50 (36%) respondents who said NO, indicating that they did not take alcohol.

## DISCUSSION

On Sex, more respondents (58%) were Females, an indication that age-related Macular Degeneration affects Females more.

On tribe, the majority (54%) of the respondents were Basoga, an indication that it is the dominant tribe in the area. On religion, most of the respondents (44%) were Catholics, while a few respondents (18%) were Muslims.

The study findings for educational levels established that the majority of the respondents (64%) were primary school dropouts, followed by 32% secondary school dropouts, while the least number of respondents had achieved a tertiary level of education. This could be because illiterates usually have poor health-seeking Behaviors.

In terms of occupation, the majority of the respondents (42%) were Peasants. According to study findings on the Home area, it was established that more respondents (78%) came from Rural areas. Such people usually have limited access to specialized health care services.

## Age-related Risk Factors

According to the social demographic data of the respondents on age, (50%) of the total number of respondents were between 61-70 years, followed by (28%) were between 51-60 years, then (18%) were between 41-50 years, while the minority(4%) were between 35-40 years.

According to the research results, 02 out of 50 respondents (04%) had Juvenile macular degeneration, and their ages

were between 30 and 40 years. This could be because Juvenile Macular Degeneration vision loss may not be noticed easily until a person reaches their 30s. Similarly, (Ziyuan et al., 2019) in their research, Deep learning for automated screening and semantic segmentation of age-related and Juvenile atrophic macular degeneration identified Juvenile macular degeneration as one of the causes of blindness among children and young people.

## Social-related risk factors

The study results showed that the majority of respondents 48% strongly agreed that smokers are at risk of age-related Macular Degeneration. This is because smoking (Nicotine) decreases plasma nitrates, nitrites, and antioxidants, thus increasing oxidative stress, which leads to degeneration, including macular degeneration. Similarly, Armstrong and Mousavi (2015) observed that smoking is a risk factor associated with AMD and asserted that current smokers are exposed to two- or three-times higher risk for AMD than non-smokers.

## Conclusion

Increasing age groups of 61-70 years and 51-60 years, smoking, and low education level were significant risk factors for age-related Macular degeneration.

## Recommendations

The Government should sensitize people on the risk factors for age-related Macular degeneration. At the same time, the Government should set up more eye clinics to increase accessibility to eye care services.

### Acknowledgment

First, I want to thank the almighty God for the wisdom and peace during my studies.

I sincerely thank my supervisor, Mr. Mulumba Taddeo, for his professional guidance and tireless effort in assisting me during the course of my study.

My appreciation also goes to Jinja Regional Referral Hospital, especially the Eye Department.

I am equally grateful to the Christian Union of Ophthalmic Clinical Officers Training School, Jinja, for being a wonderful inspiration to me.

Finally, I thank the administration of the Ophthalmic Clinical Officers' Training School for their outstanding contribution to this research report.

### LIST OF ABBREVIATIONS

<b>AMD</b>	Age-related macular degeneration
<b>CNVM</b>	Choroidal Neovascular Membranes
<b>HIV</b>	Human Immunodeficiency Virus
<b>JRRH</b>	Jinja Regional Referral Hospital

### Source of funding

The study was not funded.

### Conflict of interest

The author did not declare any conflict of interest.

### Author biography

Joshua Tumwebaze is a student of a diploma in Clinical Ophthalmology at the Ophthalmic Clinical Officers' Training School, Jinja.

Taddeo Mulumba is a tutor at the Ophthalmic Clinical Officers' Training School in Jinja.

### REFERENCES

1. Armstrong RA and Mousavi M. (2015). Overview of Risk Factors for Age-Related Macular Degeneration (AMD). *J Stem Cells* 2015; 10(3):171-91. PMID: 27125062.
2. Bamini Gopinath. (2016, August 10). Health + Medicine. Retrieved from The Conversation Africa: [https://theconversation.com/explainer-](https://theconversation.com/explainer-what-is-age-related-macular-degeneration-59889#:~:text=The%20main%20symptom%20of%20age-related%20macular%20degeneration%20is,blind%20spots%20in%20the%20centre%20of%20their%20vision.)

[what-is-age-related-macular-degeneration-59889#:~:text=The%20main%20symptom%20of%20age-related%20macular%20degeneration%20is,blind%20spots%20in%20the%20centre%20of%20their%20vision.](https://theconversation.com/explainer-what-is-age-related-macular-degeneration-59889#:~:text=The%20main%20symptom%20of%20age-related%20macular%20degeneration%20is,blind%20spots%20in%20the%20centre%20of%20their%20vision.)

3. Deng Y, Qiao L, Du M, Qu C, Wan L, Li J, Huang L. (2022). Age-related macular degeneration: epidemiology, genetics, pathophysiology, diagnosis, and targeted therapy. *Genes Dis.* 2022, 9:62-79. [10.1016/j.gendis.2021.02.009](https://doi.org/10.1016/j.gendis.2021.02.009) <https://doi.org/10.1016/j.gendis.2021.02.009>
4. F Ryan Prall, Andrew A Dahl. (2023, August 28). Drugs and Diseases >Ophthalmology. Retrieved from Medscape: <https://emedicine.medscape.com/article/1226030-overview?form=fpf>
5. Pennington KL and DeAngelis MM. (2016). Epidemiology of age-related macular degeneration (AMD): Associations with Cardiovascular disease Phenotypes and lipid factors. *Eye Vis* 2016;3:34 <https://doi.org/10.1186/s40662-016-0063-5>
6. Wong WL, Su X, Li X, Cheung CM, Klein R, Cheng CY, et al. (2014). Global prevalence of age-related macular degeneration and disease burden projection for 2020 and 2040: a systematic review and meta-analysis. *Lancet Global Health.* 2014;2(2):e106-16. [https://doi.org/10.1016/S2214-109X\(13\)70145-1](https://doi.org/10.1016/S2214-109X(13)70145-1)
7. Xiayan Xu, Jing Wu, Xiaoning, Yu Yelei, Tang Xiajing, TangandXingchaoshentu (2020). Regional differences in the Global Burden of Age-Related Macular Degeneration. *BMC Public Health* Volume 20, 410(2020) <http://doi.org/10.1186/s12889-020-8445-y>
8. Ziyuan Wang, SrinivasR, Sadda, and Zhihong Hu. (2019). Deep learning for automated screening and semantic segmentation of age-related and Juvenile atrophic macular degeneration. Vol.10950 <https://doi.org/10.1117/12.2511538>

**PUBLISHED DETAILS:**

**SJC PUBLISHERS COMPANY LIMITED**



**Category: Non Government & Non profit Organisation**

**Contact: +256 775 434 261 (WhatsApp)**

**Email: [info@sjpublisher.org](mailto:info@sjpublisher.org) or [studentsjournal2020@gmail.com](mailto:studentsjournal2020@gmail.com)**

**Website: <https://sjpublisher.org>**

**Location: Scholar's Summit Nakigalala, P. O. Box 701432, Entebbe Uganda, East Africa**